

ACCEPTED  
FOR REGISTRATION

JAN 14 2014

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

Doc ID: 537463



syngenta.

**INDIVIDUAL SALE PROHIBITED  
BY LAW**

**For Control of Commensal Rats and  
House Mice.**

**Second-Generation Anticoagulant**

Kills Warfarin-Resistant Norway Rats and  
House Mice.

Norway rats and house mice may consume a  
lethal dose in one night's feeding with first dead  
rodents appearing four or five days after  
feeding begins.

*Active Ingredient:*

Brodifacoum:

3-[3-(4'-bromo-[1,1'-biphenyl]-4-yl)-1,2,3,4-  
tetrahydro-1-naphthalenyl]-4-hydroxy-  
2H-1-benzopyran-2-one . . . . .0.005%

*Other Ingredients:* . . . . .99.995%

*Total:* . . . . .100.000%

**KEEP OUT OF REACH  
OF CHILDREN.  
CAUTION**

See additional precautionary statements and  
directions for use on label.

Distribution to and sales in "consumer" stores,  
including grocery stores, drug stores,  
hardware stores, and club stores, is prohibited.

EPA Reg. No. 100-1050 EPA Est. 61282-WI-1  
Product of the United Kingdom  
Formulated in the USA

SCP 1050A-L2F 0712 4027990

**8 pounds, 4 ounces  
(150 x 0.88 oz bait packs)**

**3/32" mini-pellets  
Net Weight**

**BAR CODE # IS  
(01) 0 07 02941 41766  
LAST DIGIT IS CHECK DIGIT  
(Barcode type: UCC/EAN 128)**

PRODUCT ID.  
**41766**



# TALON®-G Rodenticide Bait Pack Mini-Pellets

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin, or on clothing.

All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

#### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco, or using the toilet and change into clean clothing.

<b>FIRST AID</b> <b>HAVE LABEL WHEN OBTAINING TREATMENT ADVICE</b>	
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center, doctor, or <b>HOT LINE NUMBER</b> immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center, doctor, or 1-800-888-8372 immediately for treatment advice.</li></ul>
<b>HOT LINE NUMBER</b> For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) Call <b>1-800-888-8372</b>	
<b>NOTE TO PHYSICIAN</b> If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K <sub>1</sub> intramuscularly or orally. Vitamin K <sub>1</sub> is antidotal at doses of 10-20 mg ( <b>not</b> mg/kg). Repeat as necessary based on monitoring of prothrombin times.	
<b>TREATMENT FOR PET POISONING</b> If animal eats bait, call veterinarian or <b>HOT LINE NUMBER</b> at once.	
<b>NOTE TO VETERINARIAN</b> <b>Anticoagulant Brodifacoum:</b> For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K <sub>1</sub> at doses of 2-5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.	

### Environmental Hazards

This product is extremely toxic to mammals and birds. Dogs, cats, and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

## TALON®-G Rodenticide Bait Pack Mini-Pellets

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**Important: Do not expose children, pets, or other nontarget animals to rodenticides. To reduce exposure:**

- Store product not in use in a location out of reach of children and pets.
- When used in residential structures, this product must be applied in tamper-resistant bait stations. All above-ground, outdoor applications must be in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

**Note:** Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

### USE RESTRICTIONS

This product may only be used to control Norway rats, roof rats, and house mice *in and within* 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food-processing facilities, industrial and commercial buildings, trash receptacles, transport vehicles (ships, trains, aircraft), docks, and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting beyond 100 feet from a structure as defined above is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait.

Do not use in sewers. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food- or feed-handling equipment, milk- or meat-handling equipment, or surfaces that come into direct contact with food. When used in USDA-inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait.

Burrow baiting with Talon-G Rodenticide Bait Pack Mini-Pellets is prohibited.

Do not sell this product in individual containers holding less than 16 pounds of bait.

### SELECTION OF TREATMENT AREAS

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

# TALON®-G Rodenticide Bait Pack Mini-Pellets

## APPLICATION DIRECTIONS

### Norway and Roof Rats

Apply 4–16 bait packs (usually at intervals of 15–30 ft.) per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

### House Mice

Apply 1 bait pack per placement. Space placements at intervals of 8–12 ft. Two bait packs may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mouse activity cease.

### Follow-Up

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead rodents and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. Where a continuous source of infestation is present, maintain permanent bait stations and replenish with census or toxic bait as needed.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### Pesticide Storage

Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.



### Pesticide Disposal

Dispose of wastes resulting from the use of this product at an approved waste disposal facility or call your local solid waste agency for alternative disposal instructions. Never place unused product down any indoor or outdoor drain.

### Container Handling

Non-refillable container. Do not reuse or refill this container. When completely empty, offer for recycling if available or dispose of empty container in a sanitary landfill, or by incineration, or by other procedures allowed by State and local authorities.

---

Talon®, the ALLIANCE FRAME   
the SYNGENTA Logo and the PURPOSE ICON   
are registered Trademarks of a Syngenta Group Company.

Bitrex® is a registered trademark of Macfarlan Smith Ltd., Edinburgh, Scotland.

©2013 Syngenta

For non-emergency (e.g. current product information) call  
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300